Clinical Quality Indicators

Patients currently on PrEP

Patients currently on PrEP

Definitions:

• **Eligibility:** Any patient who has been prescribed PrEP during the review period.

Risk Assessment

Proportion of patients who received a risk assessment regarding history of condom usage, STI, intravenous drug use, transactional sex, and Non-occupational Post-Exposure Prophylaxis (nPEP) at baseline and every 4 months afterwards.

Denominator	Numerator
• All patients	 Number of patients who received a risk assessment at least every 4 months Risk assessments include: Condom usage STI history Intravenous drug use history Transactional sex history Non-occupational Post-Exposure
	Prophylaxis (nPEP) history

HIV Testing

Proportion of patients who received an HIV test at baseline and every 4 months afterwards.

Denominator	Numerator
 All patients 	 Number of patients who received an HIV
	test at baseline and at least every 4 months

HIV Seroconversion

Proportion of patients who test positive for HIV.

Denominator	Numerator
 All patients who were tested for HIV 	 Number of patients who test positive for
	HIV while on PrEP

STI Screening

Proportion of patients who received a syphilis, gonorrhea, and chlamydia tests at baseline and every 4 months.

Denominator	Numerator
 All patients 	 Number of patients who received syphilis,
	gonorrhea, and chlamydia tests at baseline
	and at least every 4 months

Rectal Chlamydia Testing

Proportion of patients who received a rectal chlamydia test every 4 months.

D	enominator	Nu	merator
0	All patients engaging in anal sex	0	Number of patients who were tested for
			rectal chlamydia at least every 4 months

Rectal Gonorrhea Testing

Proportion of patients who received a rectal gonorrhea test every 4 months.

Denominator	Numerator
 All patients engaging in anal sex 	 Number of patients who were tested for
	rectal gonorrhea at least every 4 months

Pharyngeal Gonorrhea Testing

Proportion of patients who received a pharyngeal gonorrhea test every 4 months.

Denominator	Numerator
 All patients engaging in oral sex 	 Number of patients who were tested for pharyngeal gonorrhea at least every 4 months

Renal Function Screening

Proportion of patients whose renal function was assessed for serum creatinine at baseline and every 6 months.

Denominator	Numerator
 All patients 	 Number of patients who received a serum
	creatinine test at baseline and every 6
	months

Urinalysis

Proportion of patients who received a urinalysis at baseline and every 6 months.

Denominator	Numerator
 All patients 	 Number of patients who received a
	urinalysis at baseline and every 6 months

Adherence Assessment

Proportion of patients who received an adherence assessment every 4 months.

Denominator	Nu	merator
 All patients 	0	Number of patients who received an
		adherence assessment every 4 months

Mental Health Screening

Proportion of patients who received a baseline and annual mental health screening.

Denominator	Numerator
 All patients 	• Number of patients who received a baseline
	and annual mental health screening

Denominator	Numerator
• All patients	 Number of patients who received a baseline
	and annual substance abuse screening
Domestic Violence Screening Proportion of patients who receive	ed a baseline and annual domestic violence screening.
Proportion of patients who receive	ed a baseline and annual domestic violence screening.
C C	

Proportion of patients who seroconverted and were linked to HIV care within 5 days.

Denominator	Nu	merator
 All patients who seroconverted 	0	Number of patients who were linked to HIV
		care within 5 days of seroconversion

Patients who have not started PrEP

Patients who have not started PrEP

Eligibility: Patients who have not been prescribed PrEP but are currently enrolled in the PrEP program during the review period.

Risk Assessment

Proportion of patients who received a risk assessment regarding history of condom usage, STI, intravenous drug use, transactional sex, and Non-occupational Post-Exposure Prophylaxis (nPEP) at baseline and every 4 months afterwards.

Denominator	Numerator		
 All patients 	 Number of patients who received a risk 		
	assessment at least every 4 months		
	 Risk assessments include: 		
	 Condom usage 		
	 STI history 		
	 Intravenous drug use history 		
	 Transactional sex history 		
	 Non-occupational Post-Exposure 		
	Prophylaxis (nPEP) history		

HIV Testing

Proportion of patients who received an HIV test every 4 months.

Denomi	nator	Nu	merator
∘ All p	patients	0	Number of patients who received an HIV
			test at least every 4 months

HIV Testing: HIV Seroconversion (not on PrEP)

Proportion of patients testing positive for HIV.

Denominator	Numerator
 All patients who were tested for HIV 	 Number of patients who test positive for
	HIV

Exclusions: Patients with a known HIV infection.

Linkage to care (Seroconversions)

Proportion of patients who seroconverted and were linked to HIV care within 5 days.

Denominator	Numerator
 All patients who seroconverted 	 Number of patients who were linked to HIV
	care within 5 days of seroconversion

STI Screening

Proportion of patients who received a syphilis, gonorrhea, and chlamydia tests at baseline and every 4 months.

Denominator Numerator

 All patients 	• Number of patients who received syphilis,
	gonorrhea, and chlamydia tests at baseline
	and at least every 4 months

Rectal Chlamydia Testing

Proportion of patients who received a rectal chlamydia test every 4 months.

Denominator	Numerator
 All patients engaging in anal sex 	 Number of patients who were tested for
	rectal chlamydia at least every 4 months

Rectal Gonorrhea Testing

Proportion of patients who received a rectal gonorrhea test every 4 months.

D	enominator	Numerator
0	All patients engaging in anal sex	 Number of patients who were tested for
		rectal gonorrhea at least every 4 months

Pharyngeal Gonorrhea Testing

Proportion of patients who received a pharyngeal gonorrhea test every 4 months.

Denominator	Numerator
 All patients engaging in oral sex 	 Number of patients who were tested for pharyngeal gonorrhea at least every 4 months
	monuns

General Clinic-Specific Quality Indicators

Clinic level Indicators

Clinic Level

Eligibility: Patients are those who:

- Are in a serodiscordant relationships
- Report having condomless sex
- Had an STI within the past 6 months
- Have injected drugs within the past 6 months
- Identify as transgender and engage in high-risk sex
- Engage in transactional sex
- Have been prescribed non-occupational post-exposure prophylaxis (nPEP).

Routine Offering of PrEP

Proportion of at risk patients who were offered PrEP.

De	enominator	Nu	merator
0	All patients	0	Number of patients who were offered PrEP
			services

Patient Experience Indicators

Patient Experience

Eligibility: All patients enrolled in the PrEP program.

Patient Experience

Proportion of patients who were assessed for experience.

De	enominator	Nu	merator
0	All patients	0	Number of patients were assessed for
			experience

Stigma

Proportion of patients who reported experiences of stigma when receiving PrEP services.

Denominator	Numerator
 All patients 	 Number of patients who reported
	experiences of stigma associated with receiving PrEP services

Access to Medication

Proportion of patients reporting gaps in medication adherence.

Denominator		Numerator	
С	All patients who have been prescribed PrEP	0	Number of patients who reported gaps in
			medication adherence

Organizational Metrics

Organizational Indicators

Ongoing Staff Training

Staff receiving annual PrEP training.

D	enominator	Numerator
С	All staff providing HIV care	 Number of staff that receive annual PrEP
		training

Stigma Reduction Training

Staff receiving annual stigma reduction training.

De	enominator	Numerator
0	All staff providing HIV care	 Number of staff that receive annual stigma
		reduction training

Referral and Linkage

Proportion of patients who were linked to primary care after stopping PrEP.

De	nominator	Numerator
0	Patients that have stopped taking PrEP	 Number of patients that are linked and
	within the review period	referred to primary care
Evel	usions: Sitos with an integrated DrED and prima	any care program

Exclusions: Sites with an integrated PrEP and primary care program.

Lost to Care

Proportion of patients who were reengaged in care after missing 2 or more consecutive PrEP clinical appointments.

De	nominator	Nu	merator	
0	All PrEP patients who have missed 2 or more	0	Number of patients reengaged in care	1
	consecutive PrEP clinical appointments			

Population Indicators

exposure prophylaxis (nPEP).

Patients with STIs		
Proportion of patients who test positive for STIs.		
Denominator	Numerator	
 All patients receiving services in a PrEP 	• Number of patients who test positive for	
program	STIs	
Number of STIs		
Incidence rate of STIs for PrEP patients.		
Denominator	Numerator	
 All patients receiving services in a PrEP 	• Number of confirmed unique diagnoses of	
program	STIs	
Proportion of identified PrEP eligible patients who v		
Denominator	Numerator	
Patients who:	• Number of patients who were linked to	
• Are in a serodiscordant relationships	PrEP services	
 Report having condomless sex 		
• Had an STI within the past 6 months		
• Have injected drugs within the past 6 months		
 Identify as transgender and engage in high- 		
risk sex		
 Engage in transactional sex 		
• Have been prescribed non-occupational post-		